## FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

## Recombinant human MDK protein



g29185 ze: Dyg prm: vailable lyophilized becies: uman spression Source: coli-derived, PGEX-4T, with N-terminal GST. ological Activity: ot tested biotoxin Level: ease contact the lab for more information econstitution: econstitution: econstitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. fter reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	EFGADCKYKFENWGACDGGTGTKVRQGTLKKARYN AQCQETIRVTKPCTPKTKAKAKAKKGKGKD (79-143 aa encoded by BC011704) Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the
P μg   prm:   vailable lyophilized   becies:   man   coli-derived, PGEX-4T, with N-terminal GST.   ological Activity:   ot tested   adotoxin Level:   ease contact the lab for more information   econstitution:   constitution:   constitution with sterile water, if glycerol has no effect   orage.   iter reconstitution with sterile water, if glycerol has no effect   a subsequent experiments, it is recommended to add an	(79-143 aa encoded by BC011704) Shipping: The product is shipped at ambient temperature.
vailable lyophilized pecies: uman spression Source: coli-derived, PGEX-4T, with N-terminal GST. ological Activity: ot tested adotoxin Level: ease contact the lab for more information econstitution: constitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. for reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
eccies: uman spression Source: coli-derived, PGEX-4T, with N-terminal GST. ological Activity: ot tested adotoxin Level: ease contact the lab for more information econstitution: econstitution: constitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. for reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
uman copression Source: coli-derived, PGEX-4T, with N-terminal GST. ological Activity: ot tested ndotoxin Level: ease contact the lab for more information econstitution: constitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. for reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
coli-derived, PGEX-4T, with N-terminal GST. ological Activity: ot tested adotoxin Level: ease contact the lab for more information econstitution: econstitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. The reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
ot tested adotoxin Level: ease contact the lab for more information econstitution: econstitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. for reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
ease contact the lab for more information econstitution: econstitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. fter reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
econstitute at 0.25 µg/µl in 200 µl sterile water for short-term orage. Ter reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	The product is shipped at ambient temperature.
orage. Ter reconstitution with sterile water, if glycerol has no effect a subsequent experiments, it is recommended to add an	
Jual volume of glycerol for long-term storage (see Stability ad Storage for more details). a different concentration is needed for your purposes please djust the reconstitution volume as required (please note: the n concentration of the final solution will vary according to e volume used). Date: Centrifuge vial before opening. When reconstituting, ently pipet and wash down the sides of the vial to ensure full covery of the protein into solution.	recommended temperature (see below).
%, by SDS-PAGE with Coomassie Brilliant Blue staining.	
ne purified protein was Lyophilized from sterile PBS (58mM Na2ł ehalose and 5 % mannitol are added as protectant before lyophi	
ore for up to 12 months at -20°C to -80°C as lyophilized powder.	
nort Term Storage: ore at 2-8°C for (1-2 weeks). Ing Term Storage: iquot and store at -20°C to -80°C for up to 3 months, constitution with sterile water and addition of an equal	
blume of glycerol. Avoid repeat freeze-thaw cycles. $74_{66} Ba \rightarrow$	
	chalose and 5 % mannitol are added as protectant before lyophi ore for up to 12 months at -20°C to -80°C as lyophilized powder. Nort Term Storage: ore at 2-8°C for (1-2 weeks). ng Term Storage: iquot and store at -20°C to -80°C for up to 3 months, constitution with sterile water and addition of an equal dume of glycerol. Avoid repeat freeze-thaw cycles.

28 kDa-

20 kDa

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522)E : proteintech@ptglab.com W:(toll free in USA), or 1(312) 455-8498ptglab.com(outside USA)(add the second technology)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.