Recombinant human PD-ECGF protein



Basic Information

Catalog Number:

Ag3052

Form:

Available lyophilized

Species:

human

Expression Source:

e coli.-derived, PGEX-4T, with N-terminal GST.

Biological Activity:

Not tested

Endotoxin Level:

Please contact the lab for more information

Peptide Sequence:

MAALMTPGTGAPPAPGDFSGEGSQGLPDPSPEPKQL PELIRMKRDGGRLSEADIRGFVAAVVNGSAQGAQIG AMLMAIRLRGMDLEETSVLTQALAQSGQQLEWPEA WRQQLVDKHSTGGVGDKVSLVLAPALAACGCKVPM ISGRGLGHTGGTLDKLESIPGFNVIQSPEQMQVLLDQ AGCCIVGQSEQLVPAEGILYAARDVTATVDSLPLITA SILSKKLVEGLSALVVDVKFGGAAVFPNOEOARELAK TLVGVGASLGLRVAAALTAMDKPLGRCVGHALEVEE ALLCMDGA

(1-297 aa encoded by BC018160)

Reconstitution and Storage

Reconstitution:

Reconstitute at 0.25 $\mu g/\mu l$ in 200 μl sterile water for short-term

After reconstitution with sterile water, if glycerol has no effect

If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used).

gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.

Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).

Purity

Formulation

Stability and Storage

Storage of Reconstituted Protein

on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details).

Note: Centrifuge vial before opening. When reconstituting,

85%, by SDS-PAGE with Coomassie Brilliant Blue staining.

Store for up to 12 months at -20°C to -80°C as lyophilized powder.

The purified protein was Lyophilized from sterile PBS (58mM Na2HPO 4,17mM NaH2PO 4, 68mM NaCl, pH8.). 5 % trehalose and 5 % mannitol are added as protectant before lyophilization. The elution buffer contain 100mM GSH.

Short Term Storage: Store at 2-8°C for (1-2 weeks).

Long Term Storage:

ptglab.com

Aliquot and store at -20°C to -80°C for up to 3 months, reconstitution with sterile water and addition of an equal volume of glycerol. Avoid repeat freeze-thaw cycles.

Selected Validation Data

