# Recombinant human ARPM1 protein



## **Basic Information**

Catalog Number:

Ag33162

Form:

Available lyophilized

Species:

human

**Expression Source:** 

e coli.-derived, PET28a, with N-terminal 6\*His.

**Biological Activity:** 

Not tested

Endotoxin Level:

Please contact the lab for more information

#### Peptide Sequence:

MIKAGVAGCREPQFIYPNIIGRAKGQSRAAQGGLELC VGDQAQDWRSSLFISYPVERGLITSCEDMEIMWKHIY DYNLKLKPCDGPVLITEPALNPLANRQQITEMFFEHLG VPAFYMSIQAVLALFAAGFTTGLVLNSGAGVTQSVPI FEGYCLPHGVQQLDLAGLDLTNYLMVLMKNHGIMLLS ASDRKIVEDIKESFCYVAMNYEEEMAKKPDCL

(16-233 aa encoded by BC007289)

# Reconstitution and Storage

#### Reconstitution:

After reconstitution with sterile water, if glycerol has no effect

If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the

gently pipet and wash down the sides of the vial to ensure full

#### Shipping:

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature (see below).

# **Purity**

## **Formulation**

Stability and Storage

Storage of **Reconstituted Protein**  Reconstitute at 0.25  $\mu g/\mu l$  in 200  $\mu l$  sterile water for short-term

on subsequent experiments, it is recommended to add an equal volume of glycerol for long-term storage (see Stability and Storage for more details).

ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting,

recovery of the protein into solution.

85%, by SDS-PAGE with Coomassie Brilliant Blue staining.

The purified protein was Lyophilized from sterile PBS (58mM Na2HPO4,17mM NaH2PO4, 68mM NaCl, pH7.4). 5 % trehalose and 5 % mannitol are added as protectant before lyophilization. The elution buffer contain 300mM imidazole.

Store for up to 12 months at -20°C to -80°C as lyophilized powder.

#### **Short Term Storage:**

Store at 2-8°C for (1-2 weeks).

Long Term Storage:

ptglab.com

Aliquot and store at -20°C to -80°C for up to 3 months, reconstitution with sterile water and addition of an equal volume of glycerol. Avoid repeat freeze-thaw cycles.

# Selected Validation Data

