FOR IN VITRO RESEARCH USE ONLY. NOT FOR USE IN HUMANS OR ANIMALS.

Recombinant human NR2C2AP protein



Biological A Not tested Endotoxin L Please cont Reconstitution and Storage After reconsoin subsequive equal volum and Storage If a different adjust the re ion concent the volume	MTHSLVCPETVSRVSSVLNRNTRQFGKKHLFDQDEI CWNSDQGPSQWVTLEFPQLIRVSQLQIQFQGGFS RRGCLEGSQGTQALHKIVDFYPEDNNSLQTFPIPAA VDRLKVTFEDATDFFGRVVIYHLRVLGEKV (1-139 aa encoded by BC057837)
50 μg Form: Available by Species: human Expression e coli-deriv Biological A Not tested Endotoxin L Please cont Reconstitution and Storage After reconson subseque equal volur and Storage If a differen adjust the re ion concent the volume	RRGCLEGSQGTQALHKIVDFYPEDNNSLQTFPIPAA VDRLKVTFEDATDFFGRVVIYHLRVLGEKV (1-139 aa encoded by BC057837)
Form: Available by Species: human Expression <i>e coli</i> -deriv Biological A Not tested Endotoxin L Please cont Please cont Please cont Reconstituti Reconstitut storage. After recons on subseque equal volur and Storage If a differen adjust the re ion concent the volume	VDRLKVTFEDATDFFGRVVIYHLRVLGEKV (1-139 aa encoded by BC057837)
Available by Species: human Expression e coli-deriv Biological A Not tested Endotoxin L Please cont Reconstitution and Storage After recons on subseque equal volur and Storage If a differen adjust the re ion concent the volume	(1-139 aa encoded by BC 057837)
Species: human Expression e coli-deriv Biological A Not tested Endotoxin L Please cont Reconstitution and Storage After recons on subseque equal volur and Storage If a differen adjust the re ion concent the volume	yophilized (1-139 aa encoded by BC057837)
human Expression e coli-deriv Biological A Not tested Endotoxin L Please cont Reconstitution and Storage After recons on subseque equal volur and Storage If a differen adjust the re ion concent the volume	
Expression e coli-deriv Biological A Not tested Endotoxin L Please cont Reconstitution Reconstitution After reconson on subseque equal volur and Storage If a differen adjust the rion concent the volume	
e coli-deriv Biological A Not tested Endotoxin L Please cont Reconstitution and Storage After recons on subseque equal volur and Storage If a differen adjust the re ion concent the volume	
Reconstitution and Storage Require the storage Reconstitution and storage Reconstitution and storage Reconstitution Reconstitu	Source:
Reconstitution and Storage If a different adjust the re ion concent the volume	ved, PGEX-4T, with N-terminal GST.
Reconstitution and Storage After reconst on subsequi equal volur and Storage If a differen adjust the re ion concent the volume	Activity:
Reconstitution and Storage After recons on subseque equal volur and Storage If a differen adjust the re ion concent the volume	
Reconstitution and Storage After recons on subseque equal volur and Storage If a differen adjust the rr ion concent the volume	Level:
and Storage After reconstitute storage. After recons on subsequi- equal volur and Storage If a differen adjust the re ion concent the volume	tact the lab for more information
and Storage After reconstitute storage. After recons on subsequi- equal volur and Storage If a differen adjust the re ion concent the volume	i
and Storage storage. After recons on subseque equal volur and Storage If a differen adjust the rr ion concent the volume	
on subseque equal volur and Storage If a differen adjust the re ion concent the volume	te at 0.25 µg/µl in 200 µl sterile water for short-term The product is shipped at ambient temperature. Upon receipt, store it immediately at the
equal volur and Storage If a differen adjust the ru ion concent the volume	stitution with sterile water, if glycerol has no effect
and Storage If a differen adjust the re ion concent the volume	ient experiments, it is recommended to add an me of glycerol for long-term storage (see Stability
If a differen adjust the re ion concent the volume	e for more details).
ion concent the volume	nt concentration is needed for your purposes please
the volume	reconstitution volume as required (please note: the
	tration of the final solution will vary according to used)
	ifuge vial before opening. When reconstituting,
• • • • •	et and wash down the sides of the vial to ensure full
recovery of	f the protein into solution.
Purity 90%, by SD	DS-PAGE with Coomassie Brilliant Blue staining.
The sector	
	d protein was Lyophilized from sterile PBS (58mM Na2HPO4,17mM NaH2PO4, 68mM NaCl, pH8.). 5 % Ind 5 % mannitol are added as protectant before lyophilization. The elution buffer contain 100mM GSH.
Stability and Storage Store for up	o to 12 months at -20°C to -80°C as lyophilized powder.
Storage of Short Term	Storage:
Store at 2.8	3°C for (1-2 weeks).
Reconstituted Protein Long Terms	
	d store at -20°C to -80°C for up to 3 months, ion with sterile water and addition of an equal
	glycerol. Avoid repeat freeze-thaw cycles.
Coloreto d Validation Data	
Selected Validation Data	74 kDa→
	$74 \text{ kDa} \rightarrow 66 \text{ kDa} \rightarrow$
	$43 \text{ kDa} \longrightarrow 43 \text{ kDa}$

20 kD

 For technical support and original validation data for this product please contact:

 T: 1 (888) 4PTGLAB (1-888-478-4522)
 E : proteintech@ptglab.com W:

 (toll free in USA), or 1(312) 455-8498
 ptglab.com

 (outside USA)
 (a)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.