For Research Use Only

Biotin-conjugated PCNA Monoclonal antibody



Purification Method:

Catalog Number: Biotin-60097

1 Publications

Basic Information

Catalog Number: GenBank Accession Number: Biotin-60097 BC000491

Biotin-60097 BC000491 Protein G purification
Size: GeneID (NCBI): CloneNo.:

100ul , Concentration: 1000 µg/ml by 5111 10D10E11

Nanodrop; Full Name: Recommended Dilutions:
Source: proliferating cell nuclear antigen IHC 1:1000-1:4000

Mouse Calculated MW:

Isotype: 29 kDa/31 kDa

IgG1 Observed MW:

Immunogen Catalog Number: 36-38 kDa

AG7416

Applications

Tested Applications:

HC IHC: human gliomas tissue,

Cited Applications:

WB

Species Specificity: human, mouse, pig, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Proliferating Cell Nuclear Antigen, commonly known as PCNA, is a protein that acts as a processivity factor for DNA polymerase δ in eukaryotic cells. This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. PCNA induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. It has to be loaded onto DNA in order to be able to stimulate APEX2. PCNA protein is highly conserved during evolution; the deduced amino acid sequences of rat and human differ by only 4 of 261 amino acids. PCNA has been used as loading control for proliferating cells. The calculated molecular weight of PCNA is 29 kDa, but modified PCNA is 36kDa (PMID: 1358458).

Positive Controls:

Notable Publications

 Author
 Pubmed ID
 Journal
 Application

 Huizhen Wang
 34789720
 Cell Death Dis
 WB

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using Biotin-60097 (PCNA antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using Biotin-60097 (PCNA antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).