For Research Use Only

Biotin-conjugated CD40L/CD154 Monoclonal antibody



Catalog Number:Biotin-66502

Basic Information

Catalog Number: GenBank Accession Number:

Biotin-66502 BC071754 GeneID (NCBI):

100ul , Concentration: 2000 $\mu g/ml$ by 959 Nanodrop: **UNIPROT ID:**

P29965 Mouse Full Name: Isotype: CD40 ligand IgG2a Calculated MW: Immunogen Catalog Number: 261 aa, 29 kDa

AG24941

Applications

Tested Applications:

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Protein A purification

Recommended Dilutions:

CloneNo.:

IHC 1:50-1:500

1E6D10

Positive Controls:

IHC: human tonsillitis tissue,

Background Information

The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33kDa transmembrane homologue or 18-kDa soluble form (sCD154). CD40L is primarily expressed on activated CD4+ T cells and on a small proportion of CD8+T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. Recent studies have suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using Biotin-66502 (CD40L/CD154 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using Biotin-66502 (CD40L/CD154 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).