

For Research Use Only

FcZero-rAb® Biotin Anti-Rat CD80 (3H5) Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA65580



Basic Information

Catalog Number:

Biotin-FcA65580

Size:

100ug , 500 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

XM_017597885

GeneID (NCBI):

25408

UNIPROT ID:

G3V671

Full Name:

Cd80 molecule

Calculated MW:

36 kDa

Purification Method:

Protein A purification

CloneNo.:

3H5

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

-

Applications

Tested Applications:

FC, ELISA

Species Specificity:

rat

Positive Controls:

FC : LPS-stimulated rat splenocytes,

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B (PMID: 7545666; 12015893; 16920215).

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ unstimulated or LPS-stimulated rat splenocytes were surface stained with 0.25 ug Biotin Anti-Rat CD80 (3H5) Rabbit IgG RecAb (Biotin-FcA65580, Clone: 3H5) and Streptavidin-PE Conjugate (PE-PF00030). Cells were then stained with FcZero-rAb™ CoraLite® Plus 647 Anti-Rat CD45RA (OX-33) Rabbit IgG Recombinant Antibody (CL647-FcA65578, Clone: OX-33). Cells were not fixed.

1x10⁶ LPS-stimulated rat splenocytes were surface stained with 0.25 ug Biotin Anti-Rat CD80 (3H5) Rabbit IgG RecAb (Biotin-FcA65580, Clone: 3H5) (red) or FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue), and Streptavidin-PE Conjugate (PE-PF00030). 1x10⁶ unstimulated rat splenocytes were surface stained with Biotin Anti-Rat CD80 (3H5) Rabbit IgG RecAb (Biotin-FcA65580, Clone:

