

For Research Use Only

# FcZero-rAb™ Biotin Anti-Human CD21 (BU32) Rabbit IgG Recombinant Antibody

Catalog Number: Biotin-FcA65591



## Basic Information

Catalog Number:	Biotin-FcA65591	GenBank Accession Number:	BC136394	Purification Method:	Protein A purification
Size:	100ug , 500 ug/ml	GenelD (NCBI):	1380	CloneNo.:	BU32
Source:	Rabbit	ENSEMBL Gene ID:	ENSG00000117322	Recommended Dilutions:	FC: 0.25 ug per 10 <sup>6</sup> cells in a 100 µl suspension
Isotype:	IgG	UNIPROT ID:	P20023	Excitation/Emission maxima wavelengths:	-
		Full Name:	complement component (3d/Epstein Barr virus) receptor 2		
		Calculated MW:	1092 aa, 119 kDa		

## Applications

Tested Applications:	FC	Positive Controls:
Species Specificity:	human	FC : human PBMCs,

## Background Information

CD21, also known as complement receptor type 2 (CR2), complement C3d receptor, and Epstein-Barr virus receptor, is a transmembrane protein that contains a small cytoplasmic domain, a transmembrane region and an extracellular domain consisting of 15 tandem short consensus repeat sequences (PMID: 6230668; 2551147). It is expressed on B cells, follicular dendritic cells, thymocytes and a subset of peripheral T cells (PMID: 26119182). CD21 binds complement fragments C3d, C3dg and iC3b and acts as a receptor for the Epstein-Barr virus (PMID: 7753047). On B cells, together with CD19 and CD81, CD21 forms a complex that functions as a co-receptor to the BCR (PMID: 7542009). It is involved in B cells activation and functions.


## Storage

**Storage:**  
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.09% sodium azide, pH7.3


For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> human PBMCs were surface stained with 0.25 ug Biotin Anti-Human CD21 (BU32) Rabbit IgG RecAb (Biotin-FcA65591, Clone: BU32) or FcZero-rAb<sup>TM</sup> Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9), and Streptavidin-PE Conjugate (PE-PF00030). Cells were then stained with APC Anti-Human CD19. Cells were not fixed.



1x10<sup>6</sup> human PBMCs were surface stained with 0.25 ug Biotin Anti-Human CD21 (BU32) Rabbit IgG RecAb (Biotin-FcA65591, Clone: BU32) (red) or FcZero-rAb<sup>TM</sup> Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue), and Streptavidin-PE Conjugate (PE-PF00030). Cells were not fixed.