For Research Use Only

FcZero-rAb™ Biotin Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG Recombinant Antibody



Purification Method:

Catalog Number:Biotin-FcA65663

Basic Information

Catalog Number: GenBank Accession Number: Biotin-FcA65663 BC031711

BC031711 Protein A purification
GeneID (NCBI): CloneNo.:

100ug, 500 ug/ml 14960 M5/114.15.2

Source: UNIPROT ID: Recommended Dilutions:

Rabbit P14434 FC: 0.25 ug per 10^6 cells in a 100 µl

Isotype: Full Name: suspension

IgG histocompatibility 2, class II antigen Excitation/Emission maxima

A, alpha wavelengths:

Applications

Tested Applications:

FC, ELISA

Species Specificity:

mouse

Positive Controls:

FC: mouse splenocytes,

Background Information

Major histocompatibility complex (MHC) class II molecules are heterodimeric transmembrane glycoproteins that are expressed on the surface of antigen-presenting cells such as dendritic cells, macrophages, and B cells. The M5/114.15.2 antibody detects polymorphic determinants present on B cells, monocytes, macrophages, dendritic cells, and activated T lymphocytes from mice carrying the H-2b, H-2d, H-2q, H-2p, H-2r and H-2u haplotypes but not from mice carrying the H-2s or H-2f haplotypes (PMID: 6170707). This antibody recognizes both I-A and I-E subregion-encoded Ia glycoproteins (I-Ab, I-Ad, I-Aq, I-Ed, I-Ek, not I-Af, I-Ak, or I-As). M5/114 antibody is reported to inhibit I-A subregion-restricted T cell responses of the H-2b, H-2d, H-2q, H-2u but not the H-2f, H-2k, H-2s haplotypes (PMID: 6173436).

Storage


Storage:


Store at 2-8 °C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 0.09% sodium azide, pH7.3

Selected Validation Data





1x10^6 mouse splenocytes were surface stained with 0.25 ug Biotin Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG RecAb (Biotin-FcA65663, Clone:M5/114.15.2), or FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) and Streptavidin-PE Conjugate (PE-PF00030). Cells were co-stained with FcZero-rAb™ APC Anti-Mouse CD3e (145-2C11) Rabbit IgG Recombinant Antibody



1x10^6 mouse splenocytes were surface stained with 0.25 ug Biotin Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2) Rabbit IgG RecAb (Biotin-FcA65663, Clone:M5/114.15.2) (red), or FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue) and Streptavidin-PE Conjugate (PE-PF00030). Cells were not fixed.