

For Research Use Only

FcZero-rAb® Biotin Anti-Human Ki-67 Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA98143-2



Basic Information

Catalog Number:

Biotin-FcA98143-2

Size:

100ug , 500 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_002417

GeneID (NCBI):

4288

UNIPROT ID:

P46013

Full Name:

antigen identified by monoclonal antibody Ki-67

Calculated MW:

359 kDa

Purification Method:

Protein A purification

CloneNo.:

241499B1

Recommended Dilutions:

FC (Intra): 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

-

Applications

Tested Applications:

FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

FC (Intra) : PHA treated human PBMCs,

Background Information

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or PHA-treated human PBMCs were intracellularly stained with 0.25 ug Biotin Anti-Human Ki-67 Rabbit RecAb (Biotin-FcA98143-2, Clone:241499B1) and Streptavidin-PE Conjugate (PE-PF00030). Cells were co-stained with FcZero-rAb® APC Anti-Human CD3 (UCHT1) Rabbit Recombinant Antibody (APC-FcA65570, Clone: UCHT1). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

1x10⁶ PHA treated human PBMCs were intracellularly stained with 0.25 ug Biotin Anti-Human Ki-67 Rabbit RecAb (Biotin-FcA98143-2, Clone:241499B1) (red) or FcZero-rAb® Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue) and Streptavidin-PE Conjugate (PE-PF00030). 1x10⁶ untreated human PBMCs were intracellularly stained with 5 ul Biotin Anti-Human Ki-67 Rabbit RecAb (Biotin-

