

For Research Use Only

# FcZero-rAb® Biotin Anti-Mouse CTLA-4/CD152 Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA98147



## Basic Information

Catalog Number:

Biotin-FcA98147

Size:

100ug, 500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG0632

GenBank Accession Number:

NM\_009843.4

GeneID (NCBI):

12477

UNIPROT ID:

P09793

Full Name:

cytotoxic T-lymphocyte-associated protein 4

Calculated MW:

25 kDa

Purification Method:

Protein A purification

CloneNo.:

241100B8

Recommended Dilutions:

FC: 0.25 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC : Con A treated mouse splenocytes,

## Background Information

CTLA-4, also known as CD152, belonging to the immunoglobulin superfamily, is primarily found on activated T cells and regulatory T cells (Tregs). CTLA-4 is closely related to the T-cell costimulatory CD28, and both molecules bind to B7-1 and B7-2 on antigen-presenting cells. CTLA-4 acts as a negative regulatory molecule of T-cell responses. Besides the full-length transmembrane form, CTLA-4 also exists in a truncated soluble form (sCTLA-4).

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> untreated or Con A-treated mouse splenocytes were surface stained with 0.25 ug Biotin Anti-Mouse CTLA-4/CD152 Rabbit RecAb (Biotin-FcA98147, Clone: 241100B8) and Streptavidin-PE Conjugate (PE-PF00030). Cells were co-stained with FcZero-rAb® APC Anti-Mouse CD4 (RM4-4) Rabbit Recombinant Antibody (APC-FcA65655, Clone: RM4-4). Cells were not fixed.

1x10<sup>6</sup> Con A-treated mouse splenocytes were surface stained with 0.25 ug Biotin Anti-Mouse CTLA-4/CD152 Rabbit RecAb (Biotin-FcA98147, Clone: 241100B8) (red) or FcZero-rAb® Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue), and Streptavidin-PE Conjugate (PE-PF00030). 1x10<sup>6</sup> untreated mouse splenocytes were surface stained with 0.25 ug Biotin Anti-Mouse CTLA-4/CD152 Rabbit RecAb (Biotin-

