

For Research Use Only

FcZero-rAb® Biotin Anti-Human CCL20/MIP-3 alpha Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA98221



Basic Information

Catalog Number:

Biotin-FcA98221

Size:

100ug , 500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG2398

GenBank Accession Number:

NM_004591.3

GeneID (NCBI):

6364

UNIPROT ID:

P78556-1

Full Name:

chemokine (C-C motif) ligand 20

Calculated MW:

11kDa

Purification Method:

Protein A purification

CloneNo.:

241694B2

Recommended Dilutions:

FC (Intra): 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

-

Applications

Tested Applications:

FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

FC (Intra) : PMA, LPS and Brefeldin A treated THP-1 cells,

Background Information

CCL20, also known as macrophage infiltrating factor protein 3a, is a C-C chemokine that specifically binds to the receptor CCR6. CCL20 is produced in various tissues and by a wide range of immune cells, including neutrophils, natural killer cells, T-helper type 17 (Th17) cells, B cells, and various antigen-presenting cells, such as dendritic cells (DCs), Langerhans cells, and macrophages. Increased expression of CCL20 is likely involved in the increased recruitment of dendritic cells observed in fibroinflammatory diseases such as chronic obstructive pulmonary disease (COPD). CCL20 expression is increased by the proinflammatory cytokine IL-1 beta.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ untreated or PMA, LPS and Brefeldin A treated THP-1 cells were surface stained with 0.25 ug Biotin Anti-Human CCL20/MIP-3 alpha Rabbit RecAb (Biotin-FcA98221, Clone: 241694B2) and PE Streptavidin (Cat.No. PF00021). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.

1x10⁶ PMA, LPS and Brefeldin A treated THP-1 cells were surface stained with 0.25 ug Biotin Anti-Human CCL20/MIP-3 alpha Rabbit RecAb (Biotin-FcA98221, Clone: 241694B2) (red) or 0.25 ug FcZero-rAb™ Biotin Rabbit IgG Isotype Control Recombinant Antibody (Biotin-FcA98136, Clone: 240953C9) (blue). 1x10⁶ untreated cells were surface stained with 0.25 ug Biotin Anti-Human CCL20/MIP-3 alpha Rabbit RecAb (Biotin-FcA98221, Clone: 241694B2)

