

For Research Use Only

FcZero-rAb™ Biotin Anti-6*His, His-Tag Rabbit Recombinant Antibody

Catalog Number: Biotin-FcA98450



Basic Information

Catalog Number:

Biotin-FcA98450

Size:

100ug, 500 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

5

UNIPROT ID:

HISTAG

Full Name:

Calculated MW:

1 kDa

Purification Method:

Protein A purification

CloneNo.:

251107A1

Recommended Dilutions:

FC (Intra): 0.25 ug per 10⁶ cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

Applications

Tested Applications:

FC (Intra), ELISA

Species Specificity:

recombinant protein

Positive Controls:

FC (Intra): Transfected HEK-293T cells,

Background Information

Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. His-tag is often used for affinity purification and binding assays. Expressed His-tagged proteins can be purified and detected easily because the string of histidine residues binds to several types of immobilized metal ions, including nickel, cobalt and copper, under specific buffer conditions. The His-tag antibody is a useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells.

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 untransfected HEK-293T cells or transfected HEK-293T cells were intracellularly stained with 0.25 μ g Biotin Anti-His tag Rabbit RecAb (Biotin-FcA98450, Clone: 251107A1), and APC-conjugated streptavidin. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.