## For Research Use Only

## CoraLite® Plus 405 Anti-Human CD8a (OKT8)

Antibodies | ELISA kits | Proteins www.ptglab.com


Catalog Number: CL405-65135

Basic Information	Catalog Number: CL405-65135	GenBank Accession Number: BC025715	Purification Method: Affinity purification
	Size: 100tests , 5 µl/test	GenelD (NCBI): 925	CloneNo.: OKT8
	Source: Mouse	ENSEMBL Gene ID: ENSG00000153563	Excitation/Emission maxima wavelengths: 399 nm / 422 nm
	Isotype: IgG2a, kappa	Full Name: CD8a molecule	
		Calculated MW: 235 aa, 26 kDa	
Applications	Tested Applications: FC		
	Species Specificity: Human		
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8 $\beta$ (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8a $\beta$ heterodimer while CD8a $\alpha$ homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) $\gamma \delta$ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).		
Storage	Storage: Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide and 0.5% BSA.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10^6 human PBMCs were surface co-stained with Coralite® Plus CL647 Anti-Human CD3 and 5 ul Coralite® Plus 405 Anti-Human CD8a (CL405-65135, Clone: OKT8) or Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.