For Research Use Only

CoraLite® Plus 405 Anti-Human CD86 (BU63)




Catalog Number:CL405-65165


| CL405-65165 | GenBank Accession Number: BC040261 | Purification Method: Protein G purification | |
|---|---|--|--|
| Size: 100tests , 5 µl/test | GenelD (NCBI): 942 | CloneNo.: BU63 | |
| Source: Mouse | ENSG00000114013 | Excitation/Emission maxima wavelengths: 399 nm / 422 nm | |
| lsotype: IgG1, kappa | | | |
| | | | |
| | Calculated MW: 329 aa, 38 kDa | | |
| Tested Applications: FC | | | |
| Species Specificity: Human | | | |
| expressed on antigen-preser for two proteins at the cell su of CD86 with CD28 antigen i | CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of CD86 with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of CD86 with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. | | |
| Storage Buffer: | Ç , , , , | ent. | |
| | Size: 100tests, 5 µl/test Source: Mouse Isotype: IgG1, kappa Tested Applications: FC Species Specificity: Human CD86 (also known as B7.2) is expressed on antigen-present for two proteins at the cell so of CD86 with CD28 antigen in Lymphocyte-associated prot Storage: Storage Buffer: | Size: GeneID (NCBI): 100tests, 5 µl/test 942 Source: ENSEMBL Gene ID: Mouse ENSG00000114013 Isotype: UNIPROT ID: IgG1, kappa P42081 Full Name: CD86 molecule Calculated MW: 329 aa, 38 kDa Tested Applications: FC Species Specificity: Human On CD86 (also known as B7.2) is a costimulatory molecule belonging to the expressed on antigen-presenting cells (APCs), including B cells, dendrit for two proteins at the cell surface of T cells, CD28 antigen and cytotoxi of CD86 with CD28 antigen is a costimulatory signal for activation of th lymphocyte-associated protein 4 negatively regulates T-cell activation Storage: Storage: Storage: Storage to light. Stable for one year after shipmed | |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10[^]6 human PBMCs were surface stained with 5 ul Coralite® Plus 405 Anti-Human CD86 (CL405-65165, Clone: BU63) or unstained. Cells were not fixed. Monocytes were gated.

1X10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 405 Anti-Human CD86 (CL405-65165, Clone: BU63) or unstained. Cells were not fixed. Monocytes were gated.