For Research Use Only

CoraLite® Plus 405 Anti-Rat CD8a (OX-8)




Catalog Number:CL405-65180

Catalog Number: CL405-65180	GenBank Accession Number: BC088126	Purification Method: Purified by protein-A affinity chromatography CloneNo.: OX-8
Size:	GenelD (NCBI): 24930 UNIPROT ID:	
100ug , 0.5 mg/ml		
Source:		
Mouse Isotype:	P07725 Full Name: CD8a molecule	Excitation/Emission maxima wavelengths: 399 nm / 422 nm
Tested Applications: FC		
Species Specificity:		
Rat		
CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8α or as a heterodimer of CD8α and CD8β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8αβ heterodimer while CD8αa homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γδ T cells and on NK cells (PMID: 211591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+ T cells (PMID: 1968084).		
Storage: Store at 2-8°C. Avoid exposu Storage Buffer: PBS with 0.09% sodium azid	re to light. Stable for one year after shipme	nt.
	CL405-65180 Size: 100ug, 0.5 mg/ml Source: Mouse Isotype: IgG1, kappa Tested Applications: FC Species Specificity: Rat CD8 is a transmembrane gly CD8a or as a heterodimer of majority of class I-restricted been found on some gut intra 2111591; 1831127; 8420975) activation (PMID: 8499079). I (PMID: 1968084).	CL405-65180 BC088126 Size: GeneID (NCBI): 100ug, 0.5 mg/ml 24930 Source: UNIPROT ID: Mouse P07725 Isotype: Full Name: IgG1, kappa CD8a molecule Tested Applications: FC Species Specificity: Rat CD8 is a transmembrane glycoprotein composed of two disulfide-linkee CD8 or as a heterodimer of CD8a and CD8β (PMID: 3264320; 8253791) majority of class I-restricted T cells express mostly the CD8aβ heterodi been found on some gut intraepithelial T cells, on some T cell receptor 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MH activation (PMID: 8499079). During T cell development, CD8 is required (PMID: 1968084). Storage: Storage: Storage: Storage: Storage: Storage to light. Stable for one year after shipme

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 Wistar rat splenocytes were surface stained with 0.5 ug CoraLite® Plus 405 Anti-Rat CD8a (CL405-65180, Clone: OX-8) or 0.5 ug Control Antibody. Cells were not fixed.