For Research Use Only

CoraLite® Plus 405 Anti-Human CD64 (10.1)



Purification Method:

Affinity purification

Excitation/Emission maxima

CloneNo.:

wavelengths: 399 nm / 422 nm

Catalog Number: CL405-65253

Basic Information

Catalog Number: CL405-65253

100 tests , 5 μ l/test

Source: Mouse Isotype:

IgG1, kappa

374 aa, 43 kDa

GenBank Accession Number:

BC032634 GeneID (NCBI):

2209

ENSEMBL Gene ID:

ENSG00000150337 **UNIPROT ID:** P12314 Full Name:

Fc fragment of IgG, high affinity Ia,

receptor (CD64) Calculated MW:

Applications

Tested Applications:

Species Specificity:

Background Information


Fcy receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or $inhibitory\ effects\ on\ cell\ functions\ after\ aggregation\ by\ complexed\ immunoglobulin\ G\ (IgG)\ (PMID:\ 17005690\).$ CD64, also known as Fcy RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 405 Anti-Human CD64 (CL405-65253, Clone:10.1) or unstained. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 5 ul Coralite® Plus 405 Anti-Human CD64 (CL405-65253, Clone:10.1) or unstained. Cells were not fixed. Monocytes were gated.