For Research Use Only

CoraLite® Plus 405 Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody



Catalog Number: CL 405-65572

Basic Information

Catalog Number: CL405-65572

Size:

100tests , 5 µl/test

Source: Mouse

Isotype: IgG2a

GenBank Accession Number:

BC032634 GeneID (NCBI):

2209

ENSEMBL Gene ID: ENSG00000150337

Full Name:

Fc fragment of IgG, high affinity Ia,

receptor (CD64) Calculated MW: 374 aa, 43 kDa

Purification Method:

Protein A purification

CloneNo.:

Excitation/Emission maxima

wavelengths: 399 nm / 422 nm

Applications

Tested Applications:

Species Specificity:

human

Background Information

Fcy receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or $inhibitory\ effects\ on\ cell\ functions\ after\ aggregation\ by\ complexed\ immunoglobulin\ G\ (IgG)\ (PMID:\ 17005690\).$ CD64, also known as Fcy RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.


Storage

Store at 2-8°C. Avoid exposure to light.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 human PBMCs were surface stained with CoraLite® Plus 647 Anti-Human CD11b and 5 ul CoraLite® Plus 405 Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody (CL405-65572, Clone: 10.1) (right), or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4) (left). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



CL405-65572 CD64(10.1),PB450-A

1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 405 Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody (CL405-65572, Clone: 10.1) (red), or CoraLite® Plus 405 Mouse IgG2a Isotype Control (C1.18.4) (CL405-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.