For Research Use Only

CoraLite® Plus 488-conjugated Beta Tubulin Polyclonal antibody



Catalog Number: CL488-10094

Basic Information

Catalog Number: CL488-10094

GenBank Accession Number:

BC000748

GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 10381

Source: tubulin, beta 3

Rabbit Calculated MW: Isotype: 50 kDa

IgG Observed MW: Immunogen Catalog Number: 55 kDa

Purification Method: Antigen affinity purification

Excitation/Emission maxima wavelengths:

488 nm / 515 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Background Information

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha, beta, and gamma tubulins have all been used as loading controls. Tubulin expression may vary according to resistance to

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human Beta Tubulin (CL488-10094) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).