

For Research Use Only

CoraLite® Plus 488-conjugated Cytokeratin 13 Polyclonal antibody

Catalog Number: CL488-10164



Basic Information

Catalog Number:	GenBank Accession Number:	Purification Method:
CL488-10164	BC002661	Antigen affinity purification
Size:	GeneID (NCBI):	Recommended Dilutions:
100ul , Concentration: 1000 ug/ml by Nanodrop;	3860	IF/ICC 1:200-1:800
Source:	UNIPROT ID:	Excitation/Emission maxima wavelengths:
Rabbit	P13646	493 nm / 522 nm
Isotype:	Full Name:	
IgG	keratin 13	
Immunogen Catalog Number:	Calculated MW:	
AG0217	50 kDa	

Applications

Tested Applications:	Positive Controls:
IF/ICC, FC (Intra)	IF/ICC : A431 cells,
Species Specificity:	
human, mouse	

Background Information

Keratin 13 is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Most of the type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. This type I cytokeratin is paired with keratin 4 and expressed in the suprabasal layers of non-cornified stratified epithelia. Mutations in keratin 13 gene and keratin 4 have been associated with the autosomal dominant disorder White Sponge Nevus. The type I cytokeratins are clustered in a region of chromosome 17q21.2.


Storage

Storage:
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed A431 cells using Coralite® Plus 488 Cytokeratin 13 antibody (CL488-10164) at dilution of 1:400.



1X10⁶ A431 cells were intracellularly stained with 1.6 ug Coralite® Plus 488 Anti-Human Cytokeratin 13 (CL488-10164) (red), or 1.6 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).