For Research Use Only

CoraLite® Plus 488-conjugated PAX8 Polyclonal antibody

www.ptglab.com

Catalog Number: CL488-10336

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

48 kDa

CL488-10336 BC001060 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7849 Nanodrop: **UNIPROT ID:** Q06710 Rabbit Full Name: Isotype paired box 8 IgG Calculated MW:

AG0306

Excitation/Emission maxima wavelengths: 493 nm / 522 nm

Purification Method:

IF/ICC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Applications

Tested Applications: IF/ICC, FC (Intra)

Immunogen Catalog Number:

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: SKOV-3 cells,

Background Information

PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain and a paired-type homeodomain. It is expressed during organogenesis of the thyroid gland, kidney and Mullerian $\,$ system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have been associated with Wilms tumor cells, thyroid and ovarian carcinomas. PAX8 is a useful marker in distinguishing ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous

This antibody can detect all 5 isoforms of PAX8 (31 kDa, 34 kDa, 41 kDa, 43 kDa and 48 kDa). A 58-60 kDa form has also been reported (PMID:15650356, 1049952). Proteintech's 10336-1-AP antibody has cross-reactivities with the other PAX family members.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 SKOV-3 cells were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human PAX8 (CL488-10336) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed SKOV-3 cells using CoraLite® Plus 488 PAX8 antibody (CL488-10336) at dilution of 1:200, CL594-phalloidin (red).