## For Research Use Only

## CoraLite® Plus 488-conjugated RRM1 Polyclonal antibody

Catalog Number:CL488-10526 Featured Product




| Basic Information      | Catalog Number:<br>CL488-10526   | GenBank Accession Number:<br>BC006498                              | Purification Method:<br>Antigen Affinity Purified             |
|------------------------|--|--|---|
|                        | Size:<br>100ul , Concentration: 1000 ug/ml by<br>Nanodrop;<br>Source:<br>Rabbit<br>Isotype:<br>IgG<br>Immunogen Catalog Number:<br>AG0789  | GenelD (NCBI):<br>6240   | Recommended Dilutions:<br>IF/ICC 1:50-1:500                   |
|                        |  | UNIPROT ID:<br>P23921<br>Full Name:<br>ribonucleotide reductase M1 | Excitation/Emission maxima<br>wavelengths:<br>493 nm / 522 nm |
|                        |  | Calculated MW:<br>90 kDa<br>Observed MW:                           |   |
|                        | Tested Applications:   | 90 kDa   | Controls  |
| Applications           | IF/ICC, FC (Intra)<br>Species Specificity:<br>human, mouse, rat, monkey  | Positive Controls:<br>IF/ICC : HepG2 cells,                        |   |
| Background Information | Ribonucleoside-diphosphate reductase functions as a heterodimer of a large and a small subunits in deoxyribonucleotide synthesis. RRM1 constitutes to the large subunit (R1) of ribonucleotide reductase, and it can either form heterodimer with small subunit RRM or RRM2B(PMID:16376858). RRM1 provides the precursors necessary for DNA synthesis. RRM1 can not be detected in quiescent cells, while its mRNA and protein are present throughout the cell cycle in cycling cells(PMID:8188248). Researches showed that RRM1 is involved in carcinogenesis, tumor progression, and the resistance of non-small-cell lung cancer (NSCLC) to treatment. Low level expression of RRM1 in NSCLC is associated with poor survival(PMID:17314339). |  |   |
| Storage                | Storage:<br>Store at -20°C. Avoid exposure to light. Stable for one year after shipment.<br>Storage Buffer:<br>PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.<br>Aliquoting is unnecessary for -20°C storage   |  |   |


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite® Plus 488 RRM1 antibody (CL488-10526) at dilution of 1:200.



1x10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488-conjugated RRM1 Polyclonal antibody (CL488-10526)(red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit 1gG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).