For Research Use Only

CoraLite® Plus 488-conjugated peroxiredoxin 2 Polyclonal antibody

Catalog Number:CL488-10545 Featured Product




Basic Information	Catalog Number: CL488-10545	GenBank Accession Number: BC003022	Purification Method: Antigen affinity Purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG0835	GenelD (NCBI): 7001 UNIPROT ID: P32119 Full Name: peroxiredoxin 2 Calculated MW: 22 kDa Observed MW: 22 kDa	Recommended Dilutions: IF/ICC 1:50-1:500
			Excitation/Emission maxima wavelengths: 493 nm / 522 nm
Applications	Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat	Positive Controls: IF/ICC : HepG2 cells,	
Background Information	PRDX2, also named as TSA, PRP, NKEFB and TDPX1, belongs to the ahpC/TSA family. It is known to act as an antioxidant enzyme whose main function is H2O2 reduction in cells. PRDX2 is involved in redox regulation of the cell. It reduces peroxides with reducing equivalents provided through the thioredoxin system. It may play an important role in eliminating peroxides generated during metabolism. PRDX2 might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H2O2. PRDX2 actions may be related to the expression of NFKB and IKB.(PMID:21248284)		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Coralite® Plus 488 peroxiredoxin 2 antibody (CL488-10545) at dilution of 1:200.



1x10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 488-conjugated peroxiredoxin 2 Polyclonal antibody (CL488-10545)(red), or 0.8 ug CoraLite® Plus 488conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).