For Research Use Only

## CoraLite® Plus 488-conjugated OPTN/Optineurin Polyclonal antibody



**Purification Method:** 

IF/ICC 1:50-1:500

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number:CL488-10837

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-10837 BC013876 GeneID (NCBI):

100ul , Concentration: 1000 ug/ml by 10133 Nanodrop: **UNIPROT ID:** Q96CV9

Rabbit Full Name: Isotype: optineurin IgG Calculated MW: Immunogen Catalog Number: 66 kDa

AG1272 Observed MW: 66 kDa, 70-74 kDa

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: MCF-7 cells,

## **Background Information**

OPTN, also named as FIP2, GLC1E, HIP7, HYPL and NRP, plays a neuroprotective role in the eye and optic nerve. It is probably part of the TNF-alpha signaling pathway that can shift the equilibrium toward induction of cell death. OPTN may act by regulating membrane trafficking and cellular morphogenesis via a complex that contains Rab8 and hungtingtin (HD). OPTN may constitute a cellular target for adenovirus E3 14.7, an inhibitor of TNF-alpha functions, thereby affecting cell death. Defects in OPTN are a cause of susceptibility to normal pressure glaucoma (NPG). OPTN mutated in adult-onset primary open angle glaucoma. It supports the protective role of OPTN in the trabecular meshwork. OPTN has 3 isoforms with MW 66,65 and 60 kDa (refer to UniProt). Catalog#10837-1-AP recognises 66 and 70-74 kDa band, and the additional 70-74 kDa band due to phosphorylation.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using CoraLite® Plus 488 OPTN antibody (CL488-10837) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human OPTN (CL488-10837) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).