For Research Use Only

## CoraLite® Plus 488-conjugated VDAC1/2 Polyclonal antibody

Catalog Number: CL488-10866 Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-10866 BC008482 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7416 Nanodrop: **UNIPROT ID:** 

P21796 Rabbit Full Name:

Isotype: voltage-dependent anion channel 1

IgG Calculated MW: Immunogen Catalog Number: 31 kDa

AG1144 Observed MW: 31 kDa

IF-P, FC (Intra)

human, mouse, rat

Antigen affinity purification Recommended Dilutions: IF-P 1:50-1:500

Excitation/Emission maxima

**Purification Method:** 

wavelengths: 493 nm / 522 nm

IF-P: mouse liver tissue,

Positive Controls: **Tested Applications: Applications** 

Species Specificity:

**Background Information** 

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This polyclonal antibody raised against full-length human VDAC1 protein can cross react with VDAC2.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using CoraLite® Plus 488 VDAC 1/2 antibody (CL488-10866) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488-conjugated VDAC1/2 Polyclonal antibody (CL488-10866)(red), or 0.8 ug CoraLite® Plus 488-conjugated Rabbit 1gG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).