For Research Use Only

CoraLite® Plus 488-conjugated HSP70 Polyclonal antibody



Catalog Number: CL488-10995

Basic Information

Catalog Number: GenBank Accession Number:

GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 3303

Source: heat shock 70kDa protein 1A

Rabbit Calculated MW: Isotype: 70 kDa Observed MW:

AG1446

CL488-10995

66-70 kDa

BC009322

Applications

Tested Applications:

Immunogen Catalog Number:

FC (Intra), IF

Species Specificity: human, mouse, rat

Recommended Dilutions: IF 1:50-1:500 Excitation/Emission maxima

Antigen affinity purification

Purification Method:

wavelengths: 493 nm / 522 nm

Background Information

HSPA1A, collectively known as HSP70 (also referred to HSP72), is a stress-inducible member of heat-shock protein 70 (HSP70) proteins which are highly conserved chaperons implicated in protein folding, protein refolding, protein transport, and protein targeting. Encoded by two closely linked, intronless and stress-inducible genes, HSPA1A and HSPA1B differ by only two amino acids and are believed to be fully interchangeable proteins. HSPA1A is a cytosol/nuclear protein able to translocate between cytoplasm and nucleus. Generally, HSPA1A is thought to be expressed in unstressed normal cells at low or undetectable levels. Expression of HSPA1A protein can be highly activated by various stressful stimuli. Significant up-regulation of HSPA1A has been found in various tumors. Recently it has been reported that HSPA1A can be constitutively expressed in selected cell types. HSP70 is a useful marker for exosome. This antibody well recognized the endogenous HSP70 protein in multiple cell lines. (21373891)

Positive Controls:

IF: HeLa cells,

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CoraLite® Plus 488 HSP70 antibody (CL488-10995) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human HSP70 (CL488-10995) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).