## For Research Use Only

## CoraLite® Plus 488-conjugated MEK1/2 Polyclonal antibody

Catalog Number:CL488-11049




Basic Information	Catalog Number: CL488-11049	GenBank Accession Number: BC018645	Purification Method: Antigen affinity purification		
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1481	GenelD (NCBI): 5605	Recommended Dilutions: IF/ICC 1:50-1:500		
		UNIPROT ID: P36507 Full Name: mitogen-activated protein kinase kinase 2	Excitation/Emission maxima wavelengths: 493 nm / 522 nm		
				Calculated MW: 44 kDa	
				Observed MW: 44-47 kDa	
		Applications	Tested Applications:	Positive Controls: IF/ICC : HeLa cells, SH-SY5Y cells	
			IF/ICC, FC (Intra) Species Specificity: human, mouse		
Background Information	MEK2(MAPK/ERK kinase 2) is also named as MAP2K2, MKK2, PRKMK2 and belongs to the MAP kinase kinase subfamily. MAPKK is itself dependent on Ser/Thr phosphorylation for activity catalyzed by MAP kinase kinase kinases (RAF or MEKK1). MEK1 and MEK2 are closely related, dual-specificity tyrosine/threonine protein kinases found in the Ras/Raf/MEK/ERK mitogen-activated protein kinase (MAPK) signaling pathway. 11049-1-AP can recognize both the MEK2 and MEK1.				
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage				


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free<br/>in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CoraLite® Plus 488 MEK1/2 antibody (CL488-11049) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human MEK1/2 (CL488-11049) (red), or 0.8 ug CoraLite® Plus 488conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).