For Research Use Only

## CoraLite®488-conjugated FUS/TLS Polyclonal antibody

Catalog Number: CL488-11570



**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-11570 BC026062 GeneID (NCBI):

Nanodrop: **UNIPROT ID:** P35637 Rabbit

100ul, Concentration: 1000 ug/ml by 2521

Isotype: fusion (involved in t(12;16) in malignant liposarcoma) IgG

Full Name:

Immunogen Catalog Number: Calculated MW: AG2150

75 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions:

Excitation/Emission maxima wavelengths:

491 nm / 516 nm

IF/ICC 1:50-1:500

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: HepG2 cells, HeLa cells

## **Background Information**

FUS (also named TLS and POMp75) belongs to the RRM TET family. FUS may play a role in the maintenance of genomic integrity; it binds both single-stranded and double-stranded DNA and promotes ATP-independent  $annealing\ of\ complementary\ single-stranded\ DNAs\ and\ D-loop\ formation\ in\ superhelical\ double-stranded\ DNA.\ FUS$ is also an RNA-binding protein, and its links to neurodegenerative disease proffer the intriguing possibility that altered RNA metabolism or RNA processing may underlie or contribute to neuron degeneration[PMID: 22640227]. FUS may be a cause of angiomatoid fibrous histiocytoma (AFH) and is implicated in certain forms of amyotrophic lateral sclerosis (ALS) and frontotemporal dementias (FTDs) such as frontotemporal lobar dementia with ubiquitin inclusions (FTLD-U)[PMID: 22640227]. This antibody is a rabbit polyclonal antibody raised against an internal region of human FUS. This antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




1X10^6 MCF-7 cells were intracellularly stained with 0.2 ug Coralite® 488 Anti-Human FUS/TLS (CL488-11570) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® 488 FUS/TLS antibody (CL488-11570) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite®488 FUS/TLS antibody (CL488-11570) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite®488 FUS/TLS antibody (CL488-11570) at dilution of 1:200, Coralite®594 TDP-43 (C-terminal) antibody (CL594-12892, red).