For Research Use Only

CoraLite® Plus 488-conjugated IFT81 Polyclonal antibody



Purification Method:

IF/ICC 1:50-1:500

493 nm / 522 nm

wavelengths:

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-11744 Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-11744 BC029349 GeneID (NCBI): 100ul, Concentration: 1000 ug/ml by 28981

Nanodrop: **UNIPROT ID:** Q8WYA0

Isotype: intraflagellar transport 81 homolog

(Chlamydomonas) IgG Immunogen Catalog Number: Calculated MW: 676 aa, 80 kDa AG2339 Observed MW:

75-80 kDa

Full Name:

Applications

Tested Applications:

Rabbit

Species Specificity: human, canine

Positive Controls:

IF/ICC: hTERT-RPE1 cells, MDCK cells

Background Information

Intraflagellar transport (IFT), mediated by molecular motors and IFT particles, is an important transport process that occurs in the cilium and has been shown to be essential for the assembly and maintenance of cilia and flagella in many organisms. IFT particles are multi-subunit complexes of proteins that functions to move non-membranebound particles from the cell body to the tip of cilium or flagellum, then return them to the cell body. Transport towards the ciliary tip is regulated by the IFT complex B (IFT-B), consisting of at least 15 IFT proteins, in association with kinesin motors, whereas transport from the ciliary tip back to the base is executed by a dynein motor in association with the IFT complex A (IFT-A), currently known to be composed of six IFT proteins. IFT81 is a subunit of IFT complex B.It may play a role in development of the testis and spermatogenesis. There are some isoforms of IFT81 with 73-78 kDa and 43-50 kDa.

Storage


Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data







Immunofluorescent analysis of (4% PFA) fixed hTERT-RPE1 cells using CoraLite® Plus 488 IFT81 antibody (CL488-11744) at dilution of 1:200, acetylated Tubulin(Lys40) antibody (66200-1-lg, Clone: 7E5H8, red).



Immunofluorescent analysis of (4% PFA) fixed MDCK cells using CoraLite® Plus 488 IFT81 antibody (CL488-11744) at dilution of 1:200, acetylated Tubulin(Lys40) antibody (66200-1-lg, Clone: 7E5H8, red).