

For Research Use Only

# CoraLite® Plus 488-conjugated S100P Polyclonal antibody

Catalog Number: CL488-11803



## Basic Information

<b>Catalog Number:</b> CL488-11803	<b>GenBank Accession Number:</b> BC006819	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 6286	<b>Excitation/Emission maximum wavelengths:</b> 493 nm / 522 nm
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P25815	
<b>Isotype:</b> IgG	<b>Full Name:</b> S100 calcium binding protein P	
<b>Immunogen Catalog Number:</b> AG2379	<b>Calculated MW:</b> 95 aa, 11 kDa	

## Applications

**Tested Applications:**  
FC (Intra)

**Species Specificity:**  
human

## Background Information

S100P protein is a relatively small (95 amino acid) isoform of the S100 protein family that was first isolated from human placenta. Overexpression of S100P has been detected in several cancers such as breast, colon, prostate, pancreatic and lung carcinomas, and the protein has been functionally implicated in carcinogenic processes. S100P protein is widely expressed in both normal and neoplastic tissues. It clearly shows ectopic expression in some cancers. Based on the high expression in certain tumors, S100P could represent a potential target for novel diagnostic and therapeutic applications.

## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

**Aliquoting is unnecessary for -20°C storage**

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human S100P (CL488-11803) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).