For Research Use Only

## CoraLite® Plus 488-conjugated STAM Polyclonal antibody



**Purification Method:** 

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Catalog Number: CL488-12434

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-12434 BC030586 GeneID (NCBI): 100ul, Concentration: 1000 ug/ml by 8027

Nanodrop: **UNIPROT ID:** Source: Q92783 Rabbit Full Name:

Isotype: signal transducing adaptor molecule (SH3 domain and ITAM motif) 1 IgG

Immunogen Catalog Number: Calculated MW: AG3112 59 kDa

> Observed MW: 70 kDa

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity: human, mouse, rat

**Background Information** 

STAM, also named as STAM1, belongs to the STAM family. It is involved in intracellular signal transduction mediated by cytokines and growth factors. Upon IL-2 and GM-CSL stimulation, it plays a role in signaling leading to DNA synthesis and MYC induction. STAM may also play a role in T-cell development. It is involved in downregulation of receptor tyrosine kinase via multivesicular body (MVBs). Together with HRS, STAM forms ESCRT-0. STAM was originally identified as an adaptor protein involved in cytokine signaling.(PMID:20505072)

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human STAM (CL488-12434) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).