For Research Use Only

CoraLite® Plus 488-conjugated PYCR1 Polyclonal antibody



Purification Method:

IF/ICC 1:50-1:500

493 nm / 522 nm

wavelengths:

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-13108

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-13108 BC022244

Size: GeneID (NCBI):
100ul , Concentration: 1000 ug/ml by 5831

Nanodrop; UNIPROT ID:

Source: P32322
Rabbit Full Name:

Isotype: pyrroline-5-carboxylate reductase 1

IgGCalculated MW:Immunogen Catalog Number:319 aa, 33.8 kDaAG3764Observed MW:

33 kDa, 35 kDa

Applications

Tested Applications: IF/ICC, FC (Intra)
Species Specificity:

human

Positive Controls:

IF/ICC : HepG2 cells,

Background Information

PYCR1, also named as P5CR1, belongs to the pyrroline-5-carboxylate reductase family. It is a housekeeping enzyme that catalyzes the last step in proline biosynthesis. PYCR1 can utilize both NAD and NADP, but has higher affinity for NAD. It is involved in the cellular response to oxidative stress. Mutation in PYCR1 will cause ARCL type II(ARCL2B). Some mutation will cause DeBarsy syndrome (DBS) which is characterized by progeroid features, ophthalmological abnormalities, intrauterine growth retardation, and cutis laxa. The MW of PYCR1 is about 33-35 kDa. PYCR1 has 3 isoforms produced by alternative splicing. This antibody may have cross reaction to PYCR2 due to the high homology.

Storage

Storage:


Store at -20 °C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 PYCR1 antibody (CL488-13108) at dilution of 1:200.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human PYCR1 (CL488-13108) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).