For Research Use Only

CoraLite® Plus 488-conjugated BST2 Polyclonal antibody

www.ptglab.com

Catalog Number: CL488-13560

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: CL488-13560 BC033873

GeneID (NCBI):

100ul, Concentration: 1000 µg/ml by 684

Source: bone marrow stromal cell antigen 2

Rabbit Calculated MW: Isotype: 180 aa, 20 kDa IgG Observed MW: Immunogen Catalog Number: 30-36 kDa, 50-70 kDa

AG4430

Applications

Tested Applications:

Species Specificity:

human, mouse

Purification Method: Antigen affinity purification Recommended Dilutions:

IF 1:200-1:800 Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Background Information

BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) $through\ ubiquitination\ and\ subsequent\ degradation.\ BST2\ may\ play\ a\ role\ in\ B-cell\ activation\ in\ rheumatoid$ arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane proteins, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 is 30-36 kDa (PMID: 19196977; 21237475).

Positive Controls: IF: HepG2 cells,

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 BST2 antibody (CL488-13560) at dilution of 1:400.