## CoraLite® Plus 488-conjugated Nanog Polyclonal antibody

Catalog Number:CL488-14295 Featured Product


Basic Information	Catalog Number: CL488-14295	GenBank Accession Number: BC 160187	Purification Method: Antigen affinity purification
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG5645	GeneID (NCBI): 79923 UNIPROT ID: Q9H9S0 Full Name: Nanog homeobox Calculated MW: 35 kDa Observed MW: 35-40 kDa	Excitation/Emission maxima wavelengths: 493 nm / 522 nm
Applications	Tested Applications: FC (Intra) Species Specificity: human, mouse, rat		
Background Information	Nanog is a member of the homeobox family of DNA binding transcription factors and has been shown to maintain embryonic stem (ES) cell self-renewal independently of leukemia inhibitory factor (LIF)/Stat3. Nanog mRNA is present in pluripotent mouse and human cell lines, and absent from differentiated cells. Functionally, Nanog works together with other key pluripotent factors (Oct4, Sox2, and Lin28) to reprogram human fibroblasts and generate induced pluripotent stem (iPS) cells. These key factors form a regulatory network to support or limit each other's expression level, which maintains the properties of ES cells. Affinity purified rabbit anti-Nanog can be used to demonstrate pluripotency of ES and IPS cells. There are two kinds of variants could recognized by NANOG, one is normal form (~39kd), the other is post-translation modified form (~48kd) (21136380). Nanog exists two isoforms with molecular weight 34.4 kDa and 31.9 kDa. (PMID: 21969378)		
Storage	Storage: Store at -20°C. Avoid exposure to ligh Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin Aliquoting is unnecessary for -20°C s	n300, 0.5% BSA, pH 7.3.	nt.

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



## Selected Validation Data



1X10^6 NCCIT cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human Nanog (CL488-14295) (red), or 0.4 ug CoraLite® Plus 488conjugated Rabbit IgC control Rabbit PolyAb (CL488-30000, Clone: ) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).