For Research Use Only

## CoraLite® Plus 488-conjugated HEXIM1 Polyclonal antibody



**Purification Method:** 

wavelengths:

488 nm / 515 nm

Antigen affinity purification

Excitation/Emission maxima

Catalog Number: CL488-15676

**Featured Product** 

**Basic Information** 

Catalog Number: CL488-15676 GenBank Accession Number: BC006460

e: GeneID (NCBI):

100ul , Concentration: 1000  $\mu g/ml$  by  $\,$  10614  $\,$ 

rop; Full Name:

Source: hexamethylene bis-acetamide

Rabbit inducible 1
Isotype: Calculated MW:
IgG 41 kDa
Immunogen Catalog Number: Observed MW:

G8144 70 kDa

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity: human, mouse, rat

**Background Information** 

HEXIM1, also named CLP1, EDG1, HIS1 or MAQ1, is best known as the inhibitor of positive transcription elongation factor b, which regulates the transcription elongation of RNA polymerase II and controls 60-70% of mRNA synthesis. HEXIM1 regulates NF-kappa-B, ESR1, NR3C1 and CIITA-dependent transcriptional activity. Previous studies showed that HEXIM1 interacts with two key p53 regulators, nucleophosmin and HDM2, implying a possible connection between HEXIM1 and the p53 signaling pathway. The mainstream MW of this protein is 70 and 54 kDa. Catalog#15676-1-AP recognises both 70 and 54 kDa band.

Storage

Storage

Store at -20°C. Avoid exposure to light.


Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human HEXIM1 (CL488-15676) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).