For Research Use Only

CoraLite® Plus 488-conjugated S100A4 Polyclonal antibody



Purification Method:

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Catalog Number: CL488-16105

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: CL488-16105 BC016300

ize: GeneID (NCBI):

100ul , Concentration: 1000 µg/ml by 6275

Source: S100 calcium binding protein A4

Rabbit Calculated MW:
Isotype: 101 aa, 12 kDa
IgG Observed MW:
Immunogen Catalog Number: 10-12 kDa

AG9019

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human, mouse

Background Information

S100A4 is a member of the S100 family of calcium-binding proteins. The S100 family members have been involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100A4 is known to localize to and function in the nucleus, cytoplasm of cells, and the extracellular space. S100A4 has also been shown to be associated with tumor growth, motility, invasion, metastasis, angiogenesis, apoptosis, and chemoresistance. It is a fibroblast-specific protein associated with mesenchymal cell morphology and motility, is expressed during epithelial-mesenchymal transformations (EMT) in vivo (PMID: 9362334). It is a specific prognostic marker for renal survival in patients with IgAN (PMID: 16105038). It is also an improved marker for lung fibroblasts that could be useful for investigating the pathogenesis of pulmonary fibrosis(PMID: 15618458). Overexpression of S100A4 is correlated with a worse prognosis inpatients with various types of cancer.

Storage

Storage

Store at -20 °C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human S100A4 (CL488-16105) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).