For Research Use Only

## CoraLite® Plus 488-conjugated NLRP3 Polyclonal antibody



**Purification Method:** 

Catalog Number: CL488-19771

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-19771 NM\_001127461 Antigen affinity purification

Size: GenelD (NCBI): Recommended Dilutions: 
100ul , Concentration: 1000 ug/ml by 114548 IF/ICC 1:50-1:500

Nanodrop; UNIPROT ID: Excitation/Emission maxima

Source: Q96P20 wavelengths:
Rabbit Full Name: 493 nm / 522 nm

Isotype: NLR family, pyrin domain containing

IgG

Calculated MW: 118 kDa Observed MW: 110 kDa

**Applications** 

Tested Applications: IF/ICC, FC (Intra)
Species Specificity:

human

Positive Controls:

IF/ICC : HepG2 cells,

## **Background Information**

NALP3, also named as C1orf7, CIAS1 and PYPAF1, belongs to the NLRP family. NALP3 may function as an inducer of apoptosis. It interacts selectively with ASC and this complex may function as an upstream activator of NF-kappa-B signaling.NALP3 inhibits TNF-alpha induced activation and nuclear translocation of RELA/NF-kB p65. Also inhibits transcriptional activity of RELA. NALP3 activates caspase-1 in response to a number of triggers including bacterial or viral infection which leads to processing and release of IL1B and IL18. Defects in NLRP3 are the cause of familial cold autoinflammatory syndrome type 1 (FCAS1) which also known as familial cold urticaria. Defects in NLRP3 are a cause of Muckle-Wells syndrome (MWS) which is urticaria-deafness-amyloidosis syndrome. Defects in NLRP3 are the cause of chronic infantile neurologic cutaneous and articular syndrome (CINCA) which also known as neonatal onset multisystem inflammatory disease (NOMID). The antibody recognizes the C-term of NALP3.

Storage

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1X10^6 THP-1 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human NLRP3 (CL488-19771) (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite® Plus 488 NLRP3 antibody (CL488-19771) at dilution of 1:200.