For Research Use Only

## CoraLite® Plus 488-conjugated DICER1 Polyclonal antibody

www.ptglab.com

Catalog Number: CL488-20567

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-20567 NM 030621 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 23405 Nanodrop: **UNIPROT ID:** 

Source: Q9UPY3 Rabbit Full Name:

Isotype: dicer 1, ribonuclease type III

IgG Calculated MW:

> 219 kDa Observed MW: 220-250 kDa, 90 kDa

wavelengths: 493 nm / 522 nm

IF/ICC 1:50-1:500

**Purification Method:** 

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

**Applications** 

**Tested Applications:** 

IF/ICC

Species Specificity:

human

Positive Controls:

IF/ICC: HepG2 cells,

## **Background Information**

DICER1, also named as DICER, HERNA and KIAA0928, belongs to the helicase family and Dicer subfamily. It is required for formation of the RNA induced silencing complex (RISC). DICER1 is a component of the RISC loading complex (RLC), also known as the micro-RNA (miRNA) loading complex (miRLC), which is composed of DICER1, EIF2C2/AGO2 and TARBP2. Within the RLC/miRLC, DICER1 and TARBP2 are required to process precursor miRNAs (pre-miRNAs) to mature miRNAs and then load them onto EIF2C2/AGO2. EIF2C2/AGO2 bound to the mature miRNA constitutes the minimal RISC and may subsequently dissociate from DICER1 and TARBP2. DICER1 cleaves doublestranded RNA to produce short interfering RNAs (siRNAs) which target the selective destruction of complementary RNAs. The antibody is specific to DICER1. The calcualted molecular weight of DICER1 is 219 kDa, but modified DICER1 is about 220-250 kDa. DICER1 exists some isoforms with molecular weight 219 kDa and 93 kDa.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

in USA), or 1(312) 455-8498 (outside USA)

## Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite® Plus 488 DICER1 antibody (CL488-20567) at dilution of 1:200.