For Research Use Only

CoraLite® Plus 488-conjugated HMGA2 Polyclonal antibody

Catalog Number: CL488-20795 Featured Product



Basic Information

Catalog Number: GenBank Accession Number:

Nanodrop; UNIPROT ID:
Source: P52926
Rabbit Full Name:

100ul, Concentration: 1000 ug/ml by 8091

Isotype: high mobility group AT-hook 2

IgG Calculated MW:
Immunogen Catalog Number: 108 aa, 12 kDa
AG14588 Observed MW:
18-20 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: IF/ICC 1:200-1:800

Excitation/Emission maxima wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

IF/ICC

Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: A549 cells,

Background Information

HMGA2 belongs to the family of high mobility group with AT-hook DNA binding domain. HMGA proteins are considered architectural transcription factors; they do not have direct transcriptional activation capacity, but instead regulate gene expression by changing DNA conformation through binding to AT-rich regions in the DNA and/or direct interaction with other transcription factors (PMID: 18202751,19551524). HMGA2 is abundantly and ubiquitously expressed and plays a crucial role during embryonic development (18425117). HMGA2 promotes stem cell self-renewal and research studies have shown that decreased HMGA2 expression is associated with stem cell aging (19551524). Investigators have shown that expression levels of HMGA2 are very low in normal adult tissues, while either overexpression or rearrangement is associated with many types of cancer (PMID: 20228781). The calcualted molecular weight of HMGA2 is 12 kDa, but modified HMGA2 is about 18-20 kDa. (PMID: 18505920)

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Ruffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed A549 cells using CoraLite® Plus 488 HMGA2 antibody (CL488-20795) at dilution of 1:400.