For Research Use Only

## CoraLite® Plus 488-conjugated NR3C2 Polyclonal antibody



**Purification Method:** 

Catalog Number: CL488-21854

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number:

CL488-21854 BC111758 Antigen affinity purification GeneID (NCBI): Recommended Dilutions: IF/ICC 1:50-1:500

100ul, Concentration: 1000 ug/ml by 4306 Nanodrop:

**UNIPROT ID:** Excitation/Emission maxima P08235 wavelengths: 493 nm / 522 nm Rabbit Full Name:

Isotype: nuclear receptor subfamily 3, group C,

IgG member 2

Immunogen Catalog Number: Calculated MW: 984 aa, 107 kDa AG16410

> Observed MW: 94-110 kDa

**Applications** 

**Tested Applications:** 

Species Specificity: human, mouse

Positive Controls:

IF/ICC: HepG2 cells,

## **Background Information**

Nuclear receptor subfamily 3 group C member 2 (NR3C2), also known as Mineralocorticoid receptor (MCR or MR), is a member of the steroid/thyroid /retinoic nuclear hormone receptor superfamily that has been shown to activate gene transcription in response to aldosterone binding. Regulation of the mineral ocorticoid receptors occurs through either receptor down-regulation (negative autoregulation) or hormone-mediated upregulation (positive autoregulation). MCR association with HSP 90 appears to be required for hormone binding to MCR and subsequent MCR activation.

Storage


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® Plus 488 NR3C2 antibody (CL488-21854) at dilution of 1:200, CL594-phalloidin (red).