For Research Use Only

CoraLite® Plus 488-conjugated TCF4 Polyclonal antibody

www.ptglab.com

Catalog Number: CL488-22337

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-22337 BC125084 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 6925 Nanodrop:

UNIPROT ID: P15884 Rabbit Full Name:

Isotype: transcription factor 4 IgG Calculated MW: Immunogen Catalog Number: 671 aa, 72 kDa AG17813 Observed MW: 72 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IF-P 1:50-1:500

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

IF-P

Species Specificity: human, mouse

Positive Controls:

IF-P: mouse skeletal muscle tissue,

Background Information

Transcription factor 4 (TCF4), also known as SEF2, ITF2, E2-2, and ME2, binds to the immunoglobulin enchancer Mu-E5/KE5-motif. Involved in the initiation of neuronal differentiation. Activates transcription by binding to the E box (5'-CANNTG-3'). Binds to the E-box present in the somatostatin receptor 2 initiator element (SSTR2-INR) to activate transcription By similarity. Human TCF4 mRNA expression is particularly high in the brain. Usage of numerous 5' exons of the human TCF4 gene potentially yields in TCF4 protein isoforms with 18 different N-termini. In addition, the diversity of isoforms is increased by alternative splicing of several internal exons. Some isoforms contain a bipartite nuclear localization signal and are exclusively nuclear, whereas distribution of other isoforms relies on heterodimerization partners. Preferentially binds to either 5'-ACANNTGT-3' or 5'-CCANNTGG-3'.TCF4 has several splicing isoforms that are expressed differentially in tissues and during cancer progression (15547706, 12796377). The scopel of molecular weight of several isoforms is about 60-90 kDa (21789225).

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skeletal muscle tissue using CoraLite® Plus 488 TCF4 antibody (CL488-22337) at dilution of 1:200. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).