CoraLite® Plus 488-conjugated mTOR Polyclonal antibody

Catalog Number:CL488-28273

Featured Product





| Basic Information | Catalog Number: CL488-28273 | GenBank Accession Number: NM_004958 | Purification Method: Antigen affinity purification | |
|-------------------------------|---|---|---|-----------------------------|
| | Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG28395 | Genel D (NCBI): 2475 | Recommended Dilutions: IF/ICC 1:50-1:500 | |
| | | UNIPROT ID: P42345 | Excitation/Emission maxima wavelengths: | |
| | | Full Name: 493 nm / 522 nm FK506 binding protein 12-rapamycin associated protein 1 Calculated MW: 289 kDa | | |
| | | | | Observed MW: 250-289 kDa |
| | | Applications | Tested Applications: Positive Controls: IF/ICC, FC (Intra) | |
| Species Specificity: human | IF/ICC : HeLa cells, HepG2 cells FC (Intra) : HeLa cells, | | | |
| Background Information | MTOR, also named as FRAP1, FRAP, FRAP2 and RAPT1, belongs to the PI3/PI4-kinase family. MTOR is a Ser/Thr protein kinase that functions as an ATP and amino acid sensor to balance nutrient availability and cell growth. MTOR is kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. MTOR has a calculated molecular mass of 289 kDa, and always can be detected at about 250 kDa due to some modifications (PMID: 14578359). | | | |
| Storage | Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3 Aliquoting is unnecessary for -20°C storage | | | |

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human mTOR (CL488-28273) (red), or 0.8 ug CoraLite® Plus 488conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Coralite® Plus 488 mTOR antibody (CL488-28273) at dilution of 1:200.