For Research Use Only

## CoraLite® Plus 488-conjugated Beta Catenin Polyclonal antibody



**Purification Method:** 

IF/ICC 1:50-1:500

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-51067

Featured Product

1 Publications

**Basic Information** 

Catalog Number: CL488-51067 GenBank Accession Number:

BC058926

e: GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 1499
Nanodrop:

Nanodrop; ENSEMBL Gene ID: Source: ENSG00000168036

Rabbit UNIPROT ID: Isotype: P35222

Full Name: catenin (cadherin-associated protein),

beta 1, 88kDa Calculated MW: 781 aa, 86 kDa Observed MW: 92 kDa

**Applications** 

Tested Applications: IF/ICC, FC (Intra)

Cited Applications:

IF

IgG

Species Specificity: human, mouse, rat, pig Positive Controls:

IF/ICC : MCF-7 cells,

## Background Information

 $\beta$ -Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein.  $\beta$ -Catenin is a 92-kDa protein, originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to a-catenin and the actin cytoskeleton. Besides its essential role in the AJs,  $\beta$ -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/ $\beta$ -catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of  $\beta$ -catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Lan Yang	39437486	Int Immunopharmacol	IF

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CoraLite® Plus 488 Beta Catenin antibody (CL488-51067) at dilution of 1:200.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human Beta Catenin (CL488-51067) (red), or 0.4 ug CoraLite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).