For Research Use Only

CoraLite® Plus 488-conjugated VDAC1/Porin Polyclonal antibody

www.ptglab.com

Purification Method:

IF-P 1:50-1:500

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number:CL488-55259 Featured Product

Basic Information

Catalog Number: GenBank Accession Number:

CL488-55259 NM 003374 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 7416 Nanodrop:

UNIPROT ID: Source: P21796 Rabbit Full Name:

Isotype: voltage-dependent anion channel 1

IgG Calculated MW:

31 kDa Observed MW: 31 kDa

Applications

Tested Applications: IF-P, FC (Intra)

Species Specificity: human, mouse, rat

Positive Controls:

IF-P: human liver cancer tissue,

Background Information

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This antibody is specific to VDAC1.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using CoraLite® Plus 488 VDAC1/Porin antibody (CL488-55259) at dilution of 1:200.

1x10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite® Plus 488-conjugated VDAC1/Porin Polyclonal antibody (CL488-55259) (red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-3000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).