For Research Use Only

CoraLite® Plus 488-conjugated MYC tag Monoclonal antibody



Catalog Number: CL488-60003 1 Publications

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein G purification

CL488-60003

GeneID (NCBI):

CloneNo.:

100ul , Concentration: 1000 ug/ml by Full Name:

1A5A2

Nanodrop:

Recommended Dilutions:

Source: Mouse

Myc tag

IF/ICC 1:50-1:500

Isotype:

Excitation/Emission maxima wavelengths:

lgG1

493 nm / 522 nm

Applications

Tested Applications:

Positive Controls:

Cited Applications:

IF/ICC: Transfected HEK-293 cells,

Species Specificity: recombinant protein

Background Information

Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and the following characteristics: solubility charexpression. The c-Myc tag corresponds to amino acid residues(EQKLISEEDL) of the human c-Myc protein. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits. Myc-Tag mouse mAb detects recombinant proteins containing the Myc tag. The antibody recognizes the Myc-tag EQKLISEEDL fused to either the amino- or carboxy- terminus of targeted proteins.

Notable Publications

Author	Pubmed ID	Journal	Application
Chunlin Lin	37573425	Oncogenesis	IF


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Transfected HEK-293 cells using Coralite® Plus 488 MYC tag antibody (CL488-60003, Clone: 1A5A2) at dilution of 1:200.