For Research Use Only

## CoraLite® Plus 488-conjugated CD3 Gamma Monoclonal antibody



**Purification Method:** 

Recommended Dilutions:

Catalog Number: CL488-60347

**Basic Information** 

Catalog Number: GenBank Accession Number: CL488-60347 BC113830

-60347 BC113830 Protein A purification
Genel D (NCBI): CloneNo.:

Size: GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 µg/ml by 917 2E9G7

Nanodrop and 1000 µg/ml by Bradford<sub>Full Name</sub>: method using BSA as the standard; CD3 molecule gamma (CD3-TC

ethod using BSA as the standard; CD3g molecule, gamma (CD3-TCR IF 1:50-1:500 complex) Fycitation/Fr

Source: complex) Excitation/Emission maxima
Mouse Calculated MW: wavelengths:
Isotype: 182 aa, 20 kDa 488 nm / 515 nm
IgG2b

Immunogen Catalog Number:

AG15502

**Applications** 

**Tested Applications:** 

IF

Species Specificity:

human

**Positive Controls:** 

IF: human tonsillitis tissue,

## **Background Information**

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells. The antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

Storage

Storage:

Store at -20 °C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CL488-60347 (CD3 gamma antibody) at dilution of 1:100.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CL488-60347 (CD3 gamma antibody) at dilution of 1:100.