For Research Use Only

CoraLite® Plus 488 Anti-Mouse CD16/32 (93)



Purification Method:

Catalog Number: CL488-65057

Basic Information

Catalog Number: GenBank Accession Number: CL488-65057 BC038070

488-65057 BC038070 Affinity purification e: GeneID (NCBI): CloneNo.:

100ug , 0.5 mg/ml 14130 93

Source: UNIPROT ID: Recommended Dilutions: P08101 IF/ICC 1:50-1:500

 Isotype:
 Full Name:
 Excitation/Emission maxima

 IgG2a, lambda
 Fc receptor, IgG, low affinity IIb
 wavelengths:

493 nm / 522 nm

Applications

Tested Applications:

IF, FC

Species Specificity:

Mouse

Positive Controls:

IF/ICC: mouse splenocytes,

Background Information

CD16 (Fcy RIII) is a 50-70 kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16 mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis. CD32 (Fcy RII) is a 40 kD transmembrane glycoprotein that binds to the Fc region of IgG with low affinity. CD32 is present on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes.

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA, pH 7.3.

Selected Validation Data



1X10^6 BALB/c mouse splenocytes were surface costained with CoraLite® Plus 647 Anti-Mouse CD3 (17A2) and 0.5 ug CoraLite® Plus 488 Anti-Mouse CD16/32 (CL488-65057, Clone:93) or 0.5 ug Isotype Control. Cells were not fixed.



Immunofluorescent analysis of mouse splenocytes using CoraLite® Plus 488 CD16/32 antibody (CL488-65057, Clone: 93) at dilution of 1:100.