For Research Use Only

CoraLite® Plus 488 Anti-Mouse CD8a (53-6.7)

Catalog Number:CL488-65069

4 Publications





Basic Information	Catalog Number: CL488-65069	GenBank A BC030679	GenBank Accession Number: Purification BC030679 Affinity puri			
	Size: 100ug , 500 µg/ml	Genel D (NO 12525	CBI):	CloneNo.: 53-6.7		
	Source: Rat	UNIPROT II P01731	UNIPROT ID: P01731		Recommended Dilutions: IF/ICC 1:250-1:1000	
	Isotype: IgG2a, kappa	Full Name: CD8 antige	n, alpha chain	Excitation/Emiss wavelengths: 493 nm / 522 nm	ion maxima	
Applications	Tested Applications: IF, FC		Positive Controls:			
	Cited Applications:	IF/ICC : mouse splenocytes,				
	Species Specificity: Mouse					
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8a or as a heterodimer of CD8a and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8a β heterodimer while CD8a α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) y δ T cells and on NK cells (PMID: 211591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).					
Notable Publications	Author	Pubmed ID	Journal Ap		Application	
	Yinghui Shang	36034656	Front Chem		IF	
	Chushu Li	34341130	J Immunother Cancer		IF	
	Mengyun Peng	38228273	J Control Release		IF	
Storage	Storage: Store at 2-8°C. Avoid expo	sure to light. Stable for	one year after shipmer	nt.		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^{^6} mouse splenocytes were surface co-stained with CoraLite®594 Anti-Mouse CD3 and 0.5 ug CoraLite® Plus 488 Anti-Mouse CD8a (CL488-65069, Clone: 53-6.7) or 0.5 ug Isotype Control. Cells were not fixed.

Immunofluorescent analysis of mouse splenocytes using CoraLite® Plus 488-conjugated Anti-Mouse CD8a (CL488-65069, Clone: 53-6.7) at dilution of 1:500.