For Research Use Only

## CoraLite® Plus 488 Anti-Human IL-1 beta (2A7C6)

Catalog Number: CL488-65276



**Basic Information** 

Catalog Number: CL488-65276

GenBank Accession Number:

BC008678

GeneID (NCBI):

100ug , 0.2 mg/mL

ENSEMBL Gene ID: Source: Mouse ENSG00000125538

Full Name: Isotype:

lgG1 interleukin 1, beta Calculated MW: Immunogen Catalog Number:

HZ-1164 269 aa, 31 kDa **Purification Method:** Affinity purification

CloneNo.:

2A7C6

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity:

human

## **Background Information**

Interleukin-1 is a pro-inflammatory cytokine with multiple biological effects. The IL-1 gene family encodes three proteins: IL-1α, IL-1β and their naturally occurring inhibitor Il-1RN. Interleukin 1β (IL-1β), mainly produced by blood monocytes and tissue macrophages, has been implicated in mediating both acute and chronic inflammation. IL-1β is known to be involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis. IL- $1\beta$  is emerging as a key mediator of carcinogenesis that characterizes host-environment interactions. (PMID: 8630372; 12401481; 23704929; 24618930)


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 0.09% sodium azide and 0.5% BSA.

in USA), or 1(312) 455-8498 (outside USA)

## **Selected Validation Data**



1X10^6 LPS and Brefeldin A treated human PBMCs were surface stained with Anti-Human HLA-DR then intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human IL-1β (CL488-65276, Clone:2A7C6) or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.



1X10^6 LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human IL-1β (CL488-65276, Clone:2A7C6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human IL-1β (CL488-65276, Clone:2A7C6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).