For Research Use Only

CoraLite® Plus 488 Anti-Human M-CSF (3G12E6)



Catalog Number: CL488-65283

Basic Information

Catalog Number:

CL488-65283

100ug , 0.2 mg/mL

Source: Mouse

Isotype: lgG1

Immunogen Catalog Number:

HZ-1192

GenBank Accession Number:

BC021117 GeneID (NCBI):

1435

Full Name: colony stimulating factor 1

(macrophage)

Calculated MW:

60 kDa

Purification Method: Affinity purification

CloneNo.: 3G12E6

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

Human

Background Information

M-CSF plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes. M-CSF promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. It also plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. On the cellular level, M-CSF promotes reorganization of the actincytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration and plays a role in lipoprotein clearance (PMID: 29734839; 29679908; 28202039)


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 LPS and TNF-a then Brefeldin A treated human PBMCs were surface stained with Coralite® Plus 405 Anti-Human CD11b then intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human M-CSF (CL488-65283, Clone:3G12E6) or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated.



1X10^6 LPS and TNF-a then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human M-CSF (CL488-65283, Clone:3G12E6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated then Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human M-CSF (CL488-65283, Clone:3G12E6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit

